

## **Deep Brain Stimulation (DBS) in patients with Parkinson's disease (PD)**

### **A Scandinavia Movement Disorder Society, ScandModis consensus document April 2021**

PD-patients in advanced stage of the disease should be referred to a Movement Disorder Clinic where a comprehensive and unbiased evaluation can be made by a neurologist specialized in movement disorders with a vast experience of adjusting oral medication and in the use of deep brain stimulation (DBS), continuous subcutaneous administration of apomorphine and continuous intestinal administration of levodopa.

Background (references 3,4,5,8,9,11,13,15,17,19,20,21,22,24,25,26,27, 28)

- Patients with Parkinson's disease, who no longer can be improved by optimizing the oral/transdermal medical treatment, have shown significant benefits from treatment with DBS (Deep Brain Stimulation) in STN, or alternatively in GPi. If troublesome tremor not responding to medical treatment is present, DBS-Vim (or DBS-Zi) can be an effective treatment option.
- DBS improves cardinal motor symptoms bradykinesia, rigidity, and tremor and reduces motor fluctuations
- DBS decreases use of medication and reduces levodopa-induced dyskinesia
- DBS increases quality of life
- There is a tendency to operate patients at an earlier time of disease than before

Criteria for referral to a specialized Movement Disorder Centre for possible device-aided treatment (references 11,15,25).

- Patients with levodopa responsive Parkinson's disease
- Preferably with a duration of Parkinson's disease > 4 years
- \*\*Moderate to severe on-off motor fluctuations
- and/or
- \*\*Moderate to severe dyskinesia
- and/or
- Medically refractory \*\*moderate to severe tremor
- \*\* Defined as moderate to severe impact on quality of life
- \*Patients with tremor-dominant PD can be treated with Vim (or Zi) DBS also >75 years of age

Exclusion criteria (references 1,2,6,7,10,16,18,26)

- Dementia
- Significant medically resistant psychiatric disease (e.g. severe depression)
- Significant medical conditions with limited life expectancy
- Conditions that prevent surgery or MRI

## Patient eligible for DBS

Patient eligibility for DBS is determined at the Movement Disorder Centre after:

- Brain imaging
- Neuropsychological assessment of cognitive function and psychiatric symptoms
- Levodopa challenge test
- NB
  - Cardiac pacemaker is not a contraindication for DBS
  - It is not necessary for the patient to be awake during surgery
  - It is not always necessary to remove all hair

## Expected outcome of DBS treatment (reference 14, 22)

Expected outcome corresponds to the effect of an optimal levodopa dosage on the motor symptoms

- Significant reduction of on-off motor fluctuations
- Significant reduction of dyskinesia
- Tremor reduction
- Decreased use of medication depending on surgical target, see below
- Improved quality of life
- Levodopa unresponsive symptoms like
  - Axial symptoms e.g. postural instability
  - Freezing of gait
  - Dysarthria
  - will not improve

## Surgery in Parkinson's disease

### Target

- The subthalamic nucleus (STN) to treat the cardinal symptoms tremor, rigidity and hypokinesia and reduce motor fluctuations, substantial reduction of medication is usually obtained leading to significant reduction of dyskinesia
- The internal part of globus pallidus (GPi) is an alternative target to treat cardinal symptoms and especially dyskinesia, however, often results in much less or no reduction of medication
- The ventral intermediate nucleus of thalamus (Vim) or zona incerta (Zi) to treat tremor only
- The electrodes are implanted bilaterally and connected to a subcutaneous lead and impulse generator (IPG) usually localized beneath the clavicle
- Each electrode has several contacts and stimulation contact and parameters are adjusted by computer telemetry

## Complications

- Surgical complications
  - Intracranial hemorrhage
  - Seizures
  - Deep Venous Thrombus
  - Pulmonary Embolism
- Hardware complications
  - Infections
  - Skin erosions
  - DBS lead-migration and fractures

## Side effects (references 7,9,12, 23, 27)

- Worsening of dysarthria
- Sometimes worsening of gait and balance especially patients > 65 years of age
- Eyelid apraxia
- Dystonia
- Psychiatric symptoms (usually transient, treatable and potentially preventable)
  - Confusion
  - Psychosis
  - Depression
  - Increased risk of suicide
  - Mania
  - Apathy (can be permanent)
- Neuropsychological symptoms
  - Reduced verbal fluency (can be permanent)
  - Increased impulsivity (can be permanent)

## Patient management and follow-up

- During the first 3-6 months frequent controls in the outpatient clinic to adjust stimulation parameters and medication to obtain maximum effect of stimulation
- In patients with directional leads stimulation may if necessary be redirected to optimize effect and/or avoid side effects
- Shared control (referral neurologist and DBS centre) of symptoms and disease development and stimulation effect
- Battery replacement every 3-5 years if relevant
- Rechargeable battery available, replacement after 15 years or more
- DBS is in general a contraindication for MRI. However MRI can be performed in most cases [at the DBS centre](#) if the MRI/neurostimulator guidelines elaborated by the manufacturer are followed
- Diathermy including shortwave diathermy, microwave diathermy or therapeutic ultrasound diathermy are contraindicated
- It is safe, however, to perform diagnostic ultrasound examination in a patient with DBS
- In case of surgery in patients with DBS monopolar electrocoagulation should be avoided. Bipolar is recommended.

If a patient with DBS needs examination by ECG, EEG or EMG the DBS can be temporarily switched off during the procedure to avoid disturbance of the examination

MRI-guided focused ultrasound ablation has been introduced as an alternative method, e.g., in patients that for medical or other reasons are not eligible for DBS. Small, but permanent unilateral lesions are made by ultrasound. Thalamus-Vim has been the most common target, but promising results have also been reported after ultrasound pallidotomy or subthalamotomy in PD patients (29,30). Future randomized longitudinal studies are needed to clarify long-term efficacy, side effects and safety regarding bilateral treatment. Equipment for MRI-guided focused ultrasound ablations are to date not available in Scandinavia.

## References

1. Ardouin C et al. Bilateral subthalamic or pallidal stimulation for Parkinson's disease affects neither memory nor executive functions: a consecutive series of 62 patients. *Ann Neurol* 1999;46:217-223.
2. Castelli L et al. Chronic deep brain stimulation of the subthalamic nucleus for Parkinson's disease: effects on cognition, mood, anxiety and personality traits. *Eur Neurol* 2006;55:136-144.
3. Deuschl G, Schade-Brittinger C, Krack P et al. A randomized trial of deep-brain stimulation for Parkinson's disease. *NEJM* 2006;355:896-9084.
4. Deuschl G et al. Stimulation of the subthalamic nucleus at an earlier disease stage of Parkinson's disease: Concept and standards of the EARLYSTIM-study. *Parkinsonism Relat Disord* 2013;19(1):56-61.
5. Schuepbach WMM, Rau J, Knudsen K, Volkmann J, Krack P, Timmermann L,....and Deuschl G, for the EARLYSTIM Study Group. Neurostimulation for Parkinson's Disease with Early Motor Complications. *N Engl J Med* 2013;368:610-22.
6. Houeto JL, Mesnage V, Mallet L et al. Behavioral disorders, Parkinson's disease, and subthalamic stimulation. *JNNP* 2002;72:701-707
7. Jahanshahi M et al. The impact of deep brain stimulation on executive function in Parkinson's disease. *Brain* 2000;123:1142-1154.
8. Krack P, Batir A, Van Biercom N et al. Five-year follow-up of bilateral stimulation of the subthalamic nucleus in advanced Parkinson's disease. *NEJM* 2003;349:1925-1934
9. Castrioto A et al. Ten-year outcome of subthalamic stimulation in Parkinson's disease: a blinded evaluation. *Arch Neurol* 2011;68:1550-1556.
10. Voon V et al. A multicentre study on suicide outcomes following subthalamic stimulation for Parkinson's disease. *Brain* 2008;131:2720-2728.
11. Moro E et al. A decision tool to support appropriate referral for deep brain stimulation in Parkinson's disease. *J Neurol* 2009;256:83-88 (and references herein).
12. Rusmann H et al. Subthalamic nucleus deep brain stimulation in Parkinson's disease over age 70 years. *Neurology* 2004;63:1952-1954.
13. The deep-brain stimulation for Parkinson's disease study group. Deep-brain stimulation of the subthalamic nucleus or the pars interna of the globus pallidus in Parkinson's disease. *NEJM* 2001;345:956-963
14. Volkmann J, Albanese A, Kulisevsky J et al. Long-term effects of pallidal or subthalamic deep brain stimulation on quality of life in Parkinson's disease. *Mov Disord* 2009;24:1154-1161
15. Volkmann J, Albanese A, Antonini A, Chaudhuri KR, Clarke CE, de Bie RM, Deuschl G, Eggert K, Houeto JL, Kulisevsky J, Nyholm D, Odin P, Østergaard K, Poewe W, Pollak P, Rabey JM, Rascol O, Rizicka E, Samuel M, Speelman H, Sydow O, Vallededoriola F, van der Linden C, Oertel W. Selecting deep brain stimulation or infusion therapies in advanced Parkinson's disease: an

- evidence-based review. *J Neurol.* 2013 Nov;260(11):2701-14.16. Voon V, Kubu C, Krack P et al. Deep brain stimulation: neuropsychological and neuropsychiatric issues. *Mov Disord* 2006;21(suppl 14):S305-S326
17. Weaver FM, Follett K, Stern M et al. Bilateral deep brain stimulation vs best medical therapy for patients with advanced Parkinson disease: a randomized controlled trial. *JAMA* 2009;301:63-73
18. Witt K, Daniels C, Reiff J et al. Neuropsychological and psychiatric changes after deep brain stimulation for Parkinson's disease: a randomised, multicentre study. *Lancet Neurol* 2008;7:605-624
19. Østergaard K, Sunde N, Dupont E. Effects of bilateral stimulation of the subthalamic nucleus in patients with severe Parkinson's disease and motor fluctuations. *Mov Disord* 2002;17: 693-700
20. Østergaard K and Sunde N. Evolution of Parkinson's disease during four years of bilateral stimulation of the subthalamic nucleus. *Mov Disord* 2006;21:624-631
21. Martinez-Ramirez D et al. Update on deep brain stimulation in Parkinson's disease. *Transl Neurodegener* 2015 June 27;4:12
22. Limousin P, Foltynie T. Long-term outcomes of deep brain stimulation in Parkinson disease. *Nat Rev Neurol.* 2019;15:234-42.
23. Tripoliti E, Zrinzo L, Martinez-Torres I, et al. Effects of subthalamic stimulation on speech of consecutive patients with Parkinson disease. *Neurology* 2011; 76: 80-86
24. Macerollo A, Zrinzo L, Akram H et al. Subthalamic nucleus deep brain stimulation for Parkinson's disease: current trends and future directions. *Expert Rev Med Devices.* 2020 Apr 6:1-12,
25. Odin P, Ray Chaudhuri K, Slevin JT, Volkmann J, Dietrichs E, Martinez-Martin P, Krauss JK, Henriksen T, Katzenschlager R, Antonini A, Rascol O, Poewe W; National Steering Committees. Collective physician perspectives on non-oral medication approaches for the management of clinically relevant unresolved issues in Parkinson's disease: Consensus from an international survey and discussion program. *Parkinsonism Relat Disord.* 2015 Oct;21(10):1133-44.
26. Dietrichs E, Odin P. Algorithms for the treatment of motor problems in Parkinson's disease. *Acta Neurol Scand.* 2017;136(5):378-385
27. Bjerknes S, Toft M, Konglund AE, Pham U, Waage TR, Pedersen L, Skjelland M, Haraldsen I, Andersson S, Dietrichs E, Skogseid IM. Multiple Microelectrode Recordings in STN-DBS Surgery for Parkinson's Disease: A Randomized Study. *Mov Disord Clin Pract.* 2018 May 8;5(3):296-305. Merola A, Romagnolo A, Krishna V, et al. Current Directions in Deep Brain Stimulation for Parkinson's Disease-Directing Current to Maximize Clinical Benefit. *Neurol Ther.* 2020 Mar 9. doi: 10.1007/s40120-020-00181-9. [Epub ahead of print] Review.
29. Moosa S, Martínez-Fernández R, Elias WJ, Del Alamo M, Eisenberg HM, Fishman PS. The role of high-intensity focused ultrasound as a symptomatic treatment for Parkinson's disease. *Mov Disord.* 2019 Sep;34(9):1243-1251. Review.
30. Martínez-Fernández R, Mániz-Miró JU, Rodríguez-Rojas R, Del Álamo M, Shah BB, Hernández-Fernández F, Pineda-Pardo JA, Monje MHG, Fernández-Rodríguez B, Sperling SA, Mata-Marín D, Guida P, Alonso-Frech F, Obeso I, Gasca-Salas C, Vela-Desojo L, Elias WJ, Obeso JA. Randomized Trial of Focused Ultrasound Subthalamotomy for Parkinson's Disease. *N Engl J Med.* 2020 Dec 24;383(26):2501-2513.